Rehabilitation Protocol: Medial Epicondylitis

Patient Name:		Date:	
---------------	--	-------	--

Phase I - Maximum Protection (0 to 10 days):

- > Complete immobilization in 90° splint
- ➤ Sling for 2 weeks
- > Ice and modalities to control inflammation

Phase II - Intermediate Phase (10 days to 4 weeks):

10 Days to 2 Weeks:

- Discontinue sling at 2 weeks
- Modalities as needed for inflammation
- > Begin passive elbow and wrist range of motion in all planes as tolerated
- ➤ Begin active shoulder retraction/protraction

Weeks 2 to 4:

- Maintain program as outlined in weeks 0 to 2
- > Continue modalities to control inflammation
- Progress to active assisted elbow and wrist range of motion in all planes
- > Initiate rotator cuff and scapular strengthening without stressing the flexor/pronator mass
- ➤ Initiate terminal range of motion stretching as tolerated

Phase III - Active Range of Motion Phase (weeks 4 to 6):

- Modalities as needed
- Continue with end range stretching of the wrist flexors and extensors
- > Begin active range of motion of the elbow and wrist in all planes
- ➤ Initiate submaximal isometrics flexor/pronator mass
- Begin PREs in wrist extension and supination
- ➤ Advance rotator cuff and scapular strengthening program
- Manual resistance and PNF patterns

Phase IV - Strengthening Phase (weeks 6 to 8):

- > Continue with terminal range stretching
- Advance PREs in wrist extension and supination
- ➤ Advance rotator cuff and scapular strengthening program
- Begin pronator/flexor mass strengthening

1

Phase V - Progressive Strengthening and Proprioceptive Phase (weeks 8 to 12):

Weeks 8 to 10:

- ➤ Increase PREs throughout wrist, elbow, and shoulder complex; include flexor/pronator mass
- > Begin global upper extremity gym strengthening program, use caution with pulling and biceps exercises

Weeks 10 to 12:

Initiate closed kinetic chain strengthening

Push-up progression

Seated serratus push-ups

> Initiate plyometric program

Plyoball wall drills

Double arm rebounder drills progressing to single arm

Phase VI - Interval Throwing Program (week 12):

- > Follow-up appointment with physician
- > Initiate return to sport program per physician approval