## Rehabilitation Protocol: Anterior Cruciate Ligament (ACL) Reconstruction with Hamstrings Autograft

Name:	Date:
Diagnosis:	Date of Surgery:
Phase I (W	veeks 0-4)
• Weigh	htbearing: As tolerated with crutches (may be modified if concomitant meniscal repair/meniscal plant or articular cartilage procedure is performed)
=	ed Knee Brace:
_	Locked in full extension for ambulation and sleeping (Weeks 0-1) Unlocked for ambulation and removed while sleeping (Weeks 1-4)
• Range	e of Motion – AAROM → AROM as tolerated
_	peutic Exercises
	Quad/Hamstring sets and heel slides
0	Non-weightbearing stretch of the Gastroc/Soleus
0	Straight-Leg Raise with brace in full extension until quad strength prevents extension lag No Hamstring Stretching Until 4 Weeks Post-Op
_	Weeks 3-12)
_	htbearing: As tolerated discontinue crutch use
_	ed Knee Brace: Discontinue brace use when patient has achieved full extension with no evidence of
	sion lag
_	e of Motion - Maintain full knee extension - work on progressive knee flexion
	peutic Exercises
0	Closed chain extension exercises
0	Hamstring Stretching, Toe Raises, Balance Exercises
0	Progress to weightbearing stretch of the Gastroc/Soleus
0	Begin use of the stationary bicycle
Phase III	(Months 3-8)
• Weigh	ntbearing: Full weightbearing
<ul> <li>Range</li> </ul>	e of Motion – Full/Painless ROM
<ul><li>Thera</li></ul>	peutic Exercises
0	Begin Hamstring strengthening
0	Advance closed chain strengthening exercises, proprioception activities
0	Begin use of the Stairmaster/Elliptical
0	Can Start Straight Ahead Running at 3 months
Phase IV	(Months 8-10)
I hase iv	
0	
0	
0	Return to sports at 10-12 months
<b>Comments:</b>	
Frequency: _	times per week Duration: weeks
Signature:	Date: