Rehabilitation Protocol: ACL and MCL Reconstruction

Namo	Date:	
Name.	Date.	
Diagnosis:	Date of Surgery:	
	ASE (Weeks 0-4)	
_	t Bearing and Range of Motion	
0	0-6 weeks: toe-touch weight bearing w/ crutches	
0	- 1	
• Brace		
	Locked in full extension at all times other than PT	
	peutic Elements:	
	Modalities as needed	
0	Patella Mob; SLR's with electric stim.; co-contractions, prone hangs	
0	,	
	No abduction of hip or leg at any time.	
Carlo	No prone hangs if PCL reconstruction!!	
• Goals:		
0		
0	1 / 0	
0	Quad control	
RECOVER	Y PHASE (Weeks 5-8)	
	nt Bearing and Range of Motion:	
0	Discontinue crutches at week 6	
• Brace	Use:	
0	At all times, open to AROM; discontinue at week 8	
Thera	peutic Elements:	
0	Continue above	
0	Gentle hip abduction with no resistance below knee	
0	Wall-sits 0-45	
0	Mini-squats with support 0-45	
0	Carpet drags (not with PCL reconstruction!!)	
0	Pool therapy	
0	Treadmill walking by 8 weeks	
Goals:		
0	a/aa/ROM: 0-0-110 by 6 weeks and free by 8 weeks	
0	SLR x 30	
0	No effusion	
STRENGTHEN PHASE (Weeks 8-12)		

• Weight Bearing and Range of Motion:

o Full

- Therapeutic Elements:
 - o Continue above with increased resistance
 - o Step-downs
 - o Treadmill
 - o Stretching

0	Begin prone hangs and HSL (if PCL reconstruction)	
Goals:		
0	Walk 1-2 miles at 15 min/mile pace	
REINTEGE	RATION PHASE (Months 3-5)	
Weight Bearing and Range of Motion:		
	Full	
Brace Use:		
0	None	
_	If return to sport, fitting for custom brace by 5 months	
0	Can start jogging/running at 6 months	
-	peutic Elements:	
	Slide boards	
o o	Begin agility drills	
0	Figure 8's	
_	Gentle loops	
0	Large zig-zags	
0	Swimming	
0	Begin plyometrics at 4 months	
• Goals:		
o douis.	Treadmill (walk 1-2 miles at 10-12 min/mile pace)	
0	Return to competitive activities	
O	Return to competitive activities	
Comments:		
comments.		
Frequency: times per week		
Januarian wood		
Signatura	Data	