## **High Tibial Osteotomy and Anterior Cruciate Ligament Reconstruction Rehabilitation Program**

Name:	Date:
Diagnosis:	Date of Surgery:
Phase I -Immediate Postoperative Phase (Day 1-5)	
• Goals:	
<ul> <li>Diminish swelling/inflammation (control hen</li> </ul>	narthrosis)
<ul> <li>Diminish postoperative pain</li> </ul>	
<ul> <li>Initiate voluntary quadriceps control</li> </ul>	
<ul> <li>Independent ambulation</li> </ul>	
• Brace:	
<ul> <li>Brace for ambulation only (POD 1 to Week 6)</li> </ul>	
<ul> <li>Weight-bearing</li> </ul>	
<ul> <li>Non-weightbearing x 6 weeks</li> </ul>	
<ul> <li>Swelling/Inflammation Control</li> </ul>	
<ul> <li>Cryotherapy</li> </ul>	
<ul> <li>Compression bandages</li> </ul>	
<ul> <li>Elevation &amp; ankle pumps</li> </ul>	
<ul> <li>Range of Motion</li> </ul>	
<ul> <li>Full passive knee extension</li> </ul>	
<ul> <li>Flexion to 45° (day 1-4)</li> </ul>	
<ul> <li>Flexion to 60° (day 5)</li> </ul>	
<ul> <li>PROM and gentle AAROM only</li> </ul>	
<ul> <li>Flexibility</li> </ul>	
<ul> <li>Hamstring and calf stretches</li> </ul>	
<ul> <li>PROM/AAROM within ROM limitations</li> </ul>	
Phase II -Acute Phase (Week 2-8)	
• Goals:	
<ul> <li>Control swelling and pain</li> </ul>	
<ul> <li>Promote healing of realignment tibial tuberos</li> </ul>	sity
<ul> <li>Quadriceps strengthening</li> </ul>	
• Brace	
<ul> <li>Continue brace for ambulation only</li> </ul>	
o Discontinue brace (week 8)	
Weight-bearing	
<ul> <li>Unlock brace for ambulation week 4</li> </ul>	
<ul> <li>Non-weight bearing x 6 weeks</li> </ul>	
o Partial weight-bearing 6-8 weeks	
<ul> <li>Discontinue Crutches at Week 8</li> </ul>	
<ul> <li>Swelling/inflammation</li> </ul>	

 $\circ \quad Elevation \\$ 

- Range of motion
  - o PROM/AAROM exercises

o Continue use of cryotherapy o Compression bandage

- ROM 0-75° (week 1-3)ROM 0-90° (week 4)
- Muscle Retraining
  - Electrical muscle stimulation to quads
  - o Quad setting isometrics
  - o Straight leg raises (flexion)
  - o Hip adduction/abduction
  - o Hip extension
  - o GENTLE submaximal isometric knee extension
  - o Week 4
    - Light leg press
    - Vertical squats (no weight)
- Flexibility
  - o Continue hamstring, calf stretches

## Phase III -Subacute Phase "Motion" Phase (Week 9-11)

- Goals
  - o Gradual improvement in ROM
  - o Improve muscular strength and endurance
  - o Control forces on extension mechanism
- Weight-bearing
  - o One crutch (week 4-6)
  - o Discontinue crutch (week 6)
- Range of motion
  - o PROM 0-115°
  - o PROM 0-125°
  - o PROM 0-125/135°
- Exercises
  - o Continue electrical muscle stimulation to quadriceps
  - o Quadriceps setting isometric
  - o Hip adduction, abuction, and extension
  - Vertical squats
  - Leg press
  - o Knee extension light (0-60°)
  - o Bicycle (week 6-8)
  - o Pool program [walking, strengthening (when able)]
- Flexibility
  - Continue all stretching exercises for LE

## Phase IV -Strengthening Phase (weeks 12-16)

- Criteria to Progress to Phase IV
  - o ROM at least 0-115 degrees
  - Absence of swelling/inflammation
  - Voluntary control of quads
- Goals
  - o Gradual improvement of muscular strength
  - Functional activities/drills
- Exercises
  - o ½ vertical squats (0-60°)
  - Wall squats (0-60°)
  - Leg press

0 0 0	Forward lunges Lateral lunges Lateral step-ups Knee extension (60-0°) Hip adduction/abduction	
0	Bicycle	
0	Stairmaster	
Phase V -Return to Activity Phase		
• Criter	ia to enter Phase V	
0	Full non-painful ROM	
0	Appropriate strength level (80% or greater of contralateralleg)	
0	Satisfactory clinical exam	
<ul> <li>Goals</li> </ul>		
0	Functional return to specific drills	
<ul> <li>Exercise</li> </ul>	ses	
0	Functional drills	
0	Strengthening exercises	
0	Flexibility exercises	
<b>Comments:</b>		
Frequency: times per week Duration: weeks		
Signature:	Date:	